

# IMPACT®

## Herbicide

For postemergence weed control in all types of field corn, popcorn or sweet corn.

### Active Ingredient:

Topramezone [3-(4,5-dihydro-3-isoxazolyl)-2-methyl-4-(methylsulfonyl)phenyl] (5-hydroxy-1-methyl-1H-pyrazol-4-yl) methanone.....29.7%  
Inert Ingredients:.....70.3%  
Total:.....100.0%

(1 gallon contains 2.8 pounds of TOPRAMEZONE free acid)

**KEEP OUT OF REACH OF CHILDREN.  
CAUTION! / PRECAUCION!**

Si usted no entiende la etiqueta, busque a alguien para que se la explique a usted en detalle.  
(If you do not understand this label, find someone to explain it to you in detail).

FIRST AID	
If in eyes:	<ul style="list-style-type: none"><li>• Hold eye open and rinse slowly and gently with water for 15 - 20 minutes.</li><li>• Remove contact lenses, if present, after first 5 minutes, then continue rinsing eye.</li><li>• Call a poison control center or doctor for treatment advice.</li></ul>
If swallowed:	<ul style="list-style-type: none"><li>• Call a poison control center or doctor immediately for treatment advice.</li><li>• Have person sip a glass of water if able to swallow.</li><li>• Do not induce vomiting unless told to do so by a poison control center or doctor.</li><li>• Do not give anything by mouth to an unconscious person.</li></ul>
If on skin or clothing:	<ul style="list-style-type: none"><li>• Take off contaminated clothing.</li><li>• Rinse skin immediately with plenty of water for 15 - 20 minutes.</li><li>• Call a poison control center or doctor for treatment advice.</li></ul>
HOT LINE NUMBER	
Have the product container or label with you when calling a poison control center or doctor, or going for treatment. FOR THE FOLLOWING EMERGENCIES, PHONE 24 HOURS A DAY: Transportation: CHEMTREC.....1-800-424-9300 Other: AMVAC.....1-323-264-3910	

## PRECAUTIONARY STATEMENTS

### HAZARDS TO HUMANS AND DOMESTIC ANIMALS

**CAUTION.** Causes moderate eye irritation. Avoid contact with skin, eyes, or clothing. Harmful if swallowed or absorbed through the skin. Wash thoroughly with soap and water after handling.

### PERSONAL PROTECTIVE EQUIPMENT (PPE)

Some materials that are chemical-resistant to this product are listed below. If you want more options, follow the instructions for Category A on an EPA chemical resistant category selection chart.

Applicators and other handlers must wear:

- Long-sleeved shirt and long pants
- Chemical resistant gloves Category A
- Shoes plus socks

Discard clothing and other absorbent materials that have been drenched or heavily contaminated with this product's concentrate. Do not reuse them. Follow manufacturer's instructions for cleaning and maintaining PPE. If no such instructions for washables, use detergent and hot water. Keep and wash PPE separately from other laundry.

### User Safety Recommendations

#### Users should:

- Wash hands before eating, drinking, chewing gum, using tobacco, or using the toilet.
- Remove clothing immediately if pesticide gets inside. Then wash thoroughly and put on clean clothing.
- Remove PPE immediately after handling this product. Wash the outside of gloves before removing. As soon as possible, wash thoroughly and change into clean clothing.

### Environmental Controls Statement

When handlers use closed systems, enclosed cabs, or aircraft in a manner that meets the requirements listed in the Worker Protection Standard (WPS) for agricultural pesticides [40 CFR 170.240(d)(4-6)], the handler PPE requirements may be reduced or modified as specified in the WPS.

### Environmental Hazards

DO NOT apply directly to water, or areas where surface water is present, or to inter-tidal areas below the mean high water mark. DO NOT contaminate water when disposing of equipment wash water. DO NOT apply this product through any type of irrigation system.

Product must be used in a manner which will prevent back siphoning in wells, spills or improper disposal of excess pesticide, spray mixtures or rinsate.

## DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling. Do not apply this product in a way that will contact workers or other persons, either directly or through drift. Only protected handlers may be in the area during application. For any requirements specific to your State or Tribe, consult the agency responsible for pesticide regulation.

### Agricultural Use Requirements

Use this product only in accordance with its labeling and with the Worker Protection Standard, 40 CFR part 170. This Standard contains requirements for the protection of agricultural workers on farms, forests, nurseries, and greenhouses, and handlers of agricultural pesticides. It contains requirements for training, decontamination, notification, and emergency assistance. It also contains specific instructions and exceptions pertaining to the statements on this label about personal protective equipment (PPE), and restricted-entry interval. The requirements in this box only apply to uses of this product that are covered by the Worker Protection Standard.

Do not enter or allow worker entry into treated areas during the restricted entry interval (REI) of 12 hours.

PPE required for early entry to treated areas that is permitted under the Worker Protection Standard and that involves contact with anything that has been treated, such as plants, soil, or water, is:

- Coveralls
- Chemical-resistant gloves Category A
- Shoes plus socks

All applicable directions, restrictions, precautions and **Limited Warranty and Disclaimer** are to be followed. This labeling must be in the user's possession during application.

## STORAGE AND DISPOSAL

Do not contaminate water, food or feed by storage or disposal.

**Pesticide Storage:** Store product in original container only. Store product in a cool, dry place. Do not store this product under wet conditions. If this product has been stored where freezing temperatures have occurred, agitate or mix contents of container well before use. Avoid cross-contamination with other pesticides.

**Pesticide Disposal:** Wastes resulting from the use of this product may be disposed of on site or at an approved waste disposal facility. Improper disposal of excess pesticide, spray mix, or rinsate is a violation of federal law. If these wastes cannot be disposed of according to label instructions, contact state agency responsible for pesticide regulation or the Hazardous Waste representative at the nearest EPA Regional Office for guidance.

**Container Disposal:** Triple rinse (or equivalent). Then offer for recycling or reconditioning, or puncture and dispose of in a sanitary landfill, or by incineration, or if allowed by state and local authorities, by burning. If burned stay out of smoke. Do not reuse container.

EPA REG. NO. 5481-524

EPA EST. NO. 34704-MS-1



4100 E. Washington Blvd.  
Los Angeles, CA 90023, USA  
1-323-264-3910 • www.amvac-chemical.com

11926-3

## In Case of Emergency

In case of large-scale spillage regarding this product, call:

CHEMTREC.....1-800-424-9300  
AMVAC.....1-323-264-3910

In case of medical emergency regarding this product, call:

- Your local doctor for immediate treatment.
- Your local poison control center (hospital).
- AMVAC 1-323-264-3910

## I. GENERAL INFORMATION

**IMPACT** is a systemic postemergence herbicide for control or growth suppression of emerged broadleaf and grass weeds in field corn (grown for grain, silage or seed), popcorn (grown for ear, kernel or seed) and sweet corn (grown for ear, kernel or seed). This product may be used on conventional and herbicide resistant/tolerant corn hybrids. AMVAC has not tested all inbred lines for tolerance to **IMPACT**. Before using **IMPACT** refer to seed company recommendations for use on inbred lines of field corn, popcorn and sweet corn.

When applied as directed at the postemergence broadcast rate of 0.75 fluid ounce per acre, **IMPACT** herbicide will control or suppress the broadleaf weeds listed in Table 1 and the grass weeds listed in Table 2.

For best performance it is recommended that **IMPACT** herbicide be tank mixed with 0.25 to 1.5 pounds active ingredient of atrazine herbicide per acre. Use lower rates of atrazine to provide added burndown of emerged weeds and the higher rates for added soil residual control. Also **IMPACT** applications must include spray additives (See section III and IV of this label for details).

**Table 1. Broadleaf Weed List, Including ALS<sup>1</sup>, Glyphosate and Triazine-Resistant Biotypes (Apply at the postemergence broadcast rate of 0.75 fluid ounce per acre.)**

Annual Broadleaf Weeds	Maximum Size (Inches) <sup>2</sup>
Amaranth, Palmer	6
Amaranth, Powell	6
Burcucumber	6
Carpenterweed	6
Chickweed, Common	4
Cocklebur, Common	8
Dandelion	6 <sup>3</sup>
Galinsoga, Hairy	6
Horseweed (Marestail)	6
Jimsonweed	6
Kochia	6
Lambsquarters, Common	6
Mallow, Common	3
Mallow, Venice	3 <sup>3</sup>
Morningglory sp.	6 <sup>3</sup>
Mustard sp.	6
Nightshade, Black	6
Nightshade, Eastern Black	6
Nightshade, Hairy	6
Pigweed, Prostrate	6
Pigweed, Redroot	6
Pigweed, Smooth	6
Pigweed, Tumble	4
Prickly Lettuce	4
Ragweed, Common	6
Ragweed, Giant	8
Sheepspurge	4
Sida, Prickly	3
Smartweed, Ladysthumb	3
Smartweed, Pennsylvania	3
Sunflower, Volunteer	8
Sunflower, Wild (common)	8
Thistle, Canada	6 <sup>3</sup>
Thistle, Russian	4
Velvetleaf	8
Waterhemp, Common	6
Waterhemp, Tall	6

<sup>1</sup>ALS (acetolactate synthase) resistant weeds include those weeds resistant to the sulfonylurea (e.g., **Spirit™** and **Exceed®**), imidazolinone (e.g., **Lightning®**), or sulfonamide (e.g., **Python™**) family of herbicides.

<sup>2</sup>For best performance spray before weeds exceed the maximum stem height or vine length listed in this table.

<sup>3</sup>Growth suppression.

**Table 2. Annual Grass Weed List, Including ALS<sup>1</sup>-Resistant Biotypes (Apply at the postemergence broadcast rate of 0.75 fluid ounce per acre.)**

Annual Grass Weeds	Maximum Leaf Stage <sup>2</sup>	Maximum Size (Inches) <sup>2</sup>
Barnyardgrass	4	4
Crabgrass, Large	4	3
Crabgrass, Smooth	4	3
Cupgrass, Woolly	3 <sup>3</sup>	3 <sup>3</sup>
Foxtail, Giant	4	4
Foxtail, Green	3	3
Foxtail, Yellow	3 <sup>3</sup>	3 <sup>3</sup>
Goosegrass	4	3
Johnsongrass, Seedling	3 <sup>3</sup>	4 <sup>3</sup>

Millet, Wild-Prosopis	3	3
Panicum, Fall	3 <sup>3</sup>	3 <sup>3</sup>
Shattercane	3 <sup>3</sup>	4 <sup>3</sup>
Signalgrass, Broadleaf	3 <sup>3</sup>	3 <sup>3</sup>

<sup>1</sup>ALS-resistant weeds include those weeds resistant to the sulfonylurea (e.g., **Accent®** and **Option®**) family of herbicides.

<sup>2</sup>For best performance spray before grasses exceed the maximum leaf stage and/or height listed in this table.

<sup>3</sup>Growth suppression.

### Mode of Action

**IMPACT** is absorbed by leaves, roots, and shoots and translocated to the growing points of sensitive weeds to provide control of emerged weeds. **IMPACT** controls weeds by inhibiting carotenoid biosynthesis (HPPD inhibitor). Temperatures and moisture conditions for active plant growth are important for optimum **IMPACT** activity. **IMPACT** applications to weeds during periods of stress conditions such as cold temperatures and/or drought, may result in reduced performance.

### Herbicide Resistance

No resistance to **IMPACT** or cross-resistance to other HPPD inhibitor herbicides is known to exist. Repeated applications of a single mode of action in a weed management plan increases the probability of selecting for naturally occurring biotypes\* with less susceptibility to herbicides using that mode of action. Therefore, **IMPACT** should be tank mixed with a herbicide having a different mode of action and/or be used in a rotation with herbicides having a different mode of action. Other HPPD inhibitor herbicides include **Balance®** (isoxaflutole) and **Callisto®** (mesotrione).

\*A weed biotype is a naturally occurring individual within a given species that has a slightly different, but distinct genetic makeup from other plants.

### Crop Tolerance

**IMPACT** should be applied during favorable growing conditions for optimum crop tolerance and weed control. Crops under environmental stress are more likely to show injury from any herbicide application. Rarely, plants under these conditions treated with **IMPACT** may show some transient bleaching of the portion of the leaves intercepting the spray application. These symptoms are temporary and occur infrequently; crop growth is not affected.

### Cultivation

Avoid disturbing (e.g., cultivation) treated areas for at least 7 days following an application of **IMPACT** to allow best herbicide uptake, translocation, and weed control.

### Insecticide Information

**IMPACT** may be used sequentially or in combination with soil or foliar applied insecticides registered for use in corn.

### Cleaning Spray Equipment

To avoid injury to sensitive crops, drain and clean application equipment thoroughly using a strong detergent or commercial sprayer cleaner according to the manufacturer's directions and then triple rinse the equipment before and after applying this product.

## II. APPLICATION INSTRUCTIONS

**IMPACT** is effective for postemergence control of annual weeds in conservation or conventional-tillage production systems.

Do not apply **IMPACT** within 30 feet of native plant community.

The applicator is responsible for any loss or damage that results from spraying **IMPACT** in a manner other than recommended in this label. In addition, applicator must follow all applicable state and local regulations and ordinances in regards to spraying.

### IMPACT Application Rate and Timing:

- Do not exceed a total of 0.75 fluid ounce per treated acre per growing season.
- **IMPACT** can be applied postemergence up to 45 days prior to corn harvest.
- Apply **IMPACT** herbicide as a postemergence treatment to corn when weeds are actively growing.
- For optimal weed control apply **IMPACT** before weeds exceed labeled height.
- **IMPACT** should be applied a minimum of one hour before rainfall or overhead irrigation.

### COVERAGE

Weeds must be thoroughly covered with spray. Dense leaf canopies shelter small weeds and can prevent adequate spray coverage.

**IMPACT** applications should be made with drop nozzles if the crop canopy prevents adequate weed coverage.

### Ground Application Methods and Equipment

Uniformly apply with properly calibrated ground equipment in 10 or more gallons of water per acre. Use higher water volumes treating larger weeds and/or high density weed infestation. Flat-fan nozzle tips are recommended for postemergence applications. A spray pressure of 20 to 40 psi is recommended.

### Spray Drift

Do not apply when weather conditions may cause drift to adjacent crops and vegetation; injury may result if this occurs. To avoid spray drift from treated areas, do not make applications when wind speeds exceed 10 mph or during periods of temperature inversions.

Use of larger droplet sizes will reduce spray drift. Agriculturally approved drift-reducing additives may also be used.

### Aerial Application Methods and Equipment

Uniformly apply with properly calibrated aerial equipment in 3 or more gallons of water per acre. Adequate spray volume must be used to provide accurate and uniform distribution of spray particles over the treated area and to avoid drift of spray particles to non-target areas.

To avoid injury to sensitive crops from drift, aerial applicators must adhere to the following SPECIAL AERIAL USE DIRECTIONS AND PRECAUTIONS.

- Nozzle height above ground must be a maximum of 10 feet.
- Nozzles must be pointed towards the rear of the aircraft. The downward angle of the nozzle should not be greater than 20 degrees.
- To minimize wing-tip vortex roll, nozzles or spray boom must not be located any closer to end of wing or rotor than three-fourths the distance from the center of the aircraft.
- Use a maximum spray pressure of 40 psi.
- A buffer zone must be established between the area to be sprayed and sensitive crops.
- DO NOT spray when wind velocity is greater than 5 mph or during periods of temperature inversions. Coarse sprays (larger droplets) are less likely to drift.

### III. ADDITIVES

Postemergence applications of **IMPACT** require the addition of an adjuvant and a nitrogen fertilizer source to achieve optimum weed control.

**I. ADJUVANTS:** Unless specific recommendations are given in Section VII. Crop Use Directions - Tank Mixes, always use a methylated seed oil (MSO) or a petroleum-based or vegetable seed-based oil concentrate (COC) with **IMPACT**. For best performance across a wide range of environmental conditions, including when weeds are under moisture and/or temperature stress, use of an MSO adjuvant is recommended. Apply these oil based adjuvant concentrates at the rates of 1.0 to 1.5 gallons per 100 gallons of water (1.0% to 1.5% v/v). Use the higher rate when making an application during periods of hot dry weather.

#### AND

**II. NITROGEN FERTILIZER SOURCE:** Recommended nitrogen based fertilizers include urea ammonium nitrate (UAN; 28-34%) or ammonium phosphate (10-34-0) at 1.25 to 2.5 gallons per hundred gallons of water (1.25% to 2.5% v/v). Instead of a liquid fertilizer, spray grade ammonium sulfate (AMS) at 8.5 to 17 pounds per 100 gallons of water may be used. Use the higher rate when making application during periods of hot dry weather.

Agriculturally approved drift-reducing additives may be used in applications with **IMPACT**.

### IV. MIXING ORDER

Liquid fertilizer as a carrier for postemergence applications of **IMPACT** herbicide is not recommended. Use only water as a carrier.

#### WATER:

1. Fill the spray tank  $\frac{1}{2}$  to  $\frac{3}{4}$  full with clean water.
2. Add the required amount of **IMPACT** to the spray tank while agitating.
3. After the **IMPACT** has visibly dispersed, add spray additives and fill the remainder of the tank with water.

#### TANK-MIX PREPARATION:

When tank-mixing **IMPACT** with recommended herbicides, add the other herbicides and other components in the following order, while agitating:

1. Fill spray tank  $\frac{1}{2}$  to  $\frac{3}{4}$  full with clean water.
2. Add soluble packet products and thoroughly mix.
3. Add **IMPACT** and thoroughly mix.
4. Add WP (wetttable powder), DG (dispersible granule), DF (dry flowable), or LF (liquid flowable) formulations.
5. Add EC (emulsifiable concentrate) products.
6. Add spray adjuvants to the spray tank.
7. While agitating, fill the remainder of the tank with water.

### V. GENERAL TANK MIX INFORMATION

**IMPACT** is recommended to be used sequentially or tank mixed with other herbicides as part of a complete weed control program. Tank mix recommendations are for use only in states where the sequential or tank mix product and application site is registered. Refer to section VII. Crop Use Directions for more details and for specific tank mix restrictions. Read and follow the applicable **Restrictions and Limitations and Directions for Use** on all products included in any tank mix. **The most restrictive labeling applies to tank mixes.**

### VI. GENERAL RESTRICTIONS AND LIMITATIONS

- No more than 0.75 fluid ounce per acre (0.0164 lbs ai/acre) of **IMPACT** may be applied during the growing season.
- In the event of a crop loss due to weather, any corn type can be replanted following an application of **IMPACT** herbicide. If **IMPACT** was tank-mixed with other herbicides, the label restrictions for these herbicides must also be followed.
- DO NOT apply **IMPACT** within 45 days of corn harvest (fresh market sweet corn, silage, fodder, or grain).
- DO NOT graze or feed treated corn forage, silage, fodder, or grain for at least 45 days after an application of **IMPACT**.

**ROTATIONAL CROPS GUIDELINE:** The following rotational crops may be planted after applying **IMPACT** herbicide at the recommended rate in corn. Planting earlier than the recommended interval may result in crop injury. Avoid over applications by minimizing overlaps of spray swaths and by switching off spray boom when turning (end rows).

Minimum Time From Application to Planting	Crops
Anytime	All types of field corn, sweet corn and popcorn.
Three months	Cereal crops (wheat, barley, oats, and rye).
Nine months	Alfalfa, cotton, peanut, pea, potato, sorghum and sunflower. Soybean (except for geographic areas defined in eighteen months category).
Eighteen months	Soybean - For use in SD, ND, MN, MI and WI. All other crops not listed in categories above.

### VII. CROP USE DIRECTIONS

#### Corn (Field, Pop, Seed, and Sweet)

**IMPACT** can be applied postemergence on all corn types, including conventional, **Clearfield**®, **Roundup Ready**®, and **LibertyLink**® hybrids. In addition, **IMPACT** may be used on inbred lines used in field corn, popcorn and sweet corn seed production. Refer to seed company recommendations before use on inbred lines.

**IMPACT** may be used in tank mixtures or sequential applications with other herbicides registered for use in corn. If **IMPACT** is tank mixed with other herbicides, the label restrictions for the most restrictive of the tank mix products should be followed.

#### TANK MIXES

**Recommended tank mixes for postemergence applications of IMPACT when used at 0.75 fluid ounces per acre include:**

2,4-D <sup>1</sup>	Harness®
Aim®	Harness® Xtra
Atrazine	Hornet™
Basagran®	Keystone™
Bicep II Magnum®	Keystone™ LA
Bicep Lite II Magnum®	Laddok®
Buctril®	Marksman® <sup>1</sup>
Cinch™	Outlook™
Clarity® <sup>1</sup>	Permit®
Degree™	Prowl® H <sub>2</sub> O
Degree Xtra™	Status®
Distinct® <sup>1</sup>	Stinger™
Dual II Magnum®	Surpass™
G-Max Lite™	TopNotch™
Guardman Max™	Tough®

**Recommended tank mixes for postemergence applications of IMPACT when used at 0.5 to 0.75 fluid ounces per acre include<sup>2</sup>:**

Accent®	Roundup® brands
Glyphosate brands	Steadfast®
Liberty®	Stout™
Lightning®	Touchdown® brands
Option®	

<sup>1</sup> Use of methylated seed oil (MSO) or crop oil concentrate (COC) in tank mixes of **IMPACT** plus 2,4-D, Clarity, Distinct, or Marksman may result in crop injury if applied during periods of cold, wet weather or hot and/or humid weather. Under these environmental conditions the use of a nonionic surfactant in place of an oil based adjuvant is recommended.

<sup>2</sup> **IMPACT** may be applied at a reduced rate of 0.5 to 0.75 fluid ounce per acre to provide additional control of emerged broadleaf weeds when used in tank mixes with listed postemergence herbicides. For best performance on targeted broadleaf weeds, do not exceed the maximum labeled weed height found in Table 1 under General Information.

#### Sequential Herbicide Combinations and Uses

In addition to the control of many emerged broadleaf weeds, **IMPACT** controls or suppresses growth of several emerged grass weed species. To target a broader spectrum of grasses **IMPACT** should be used as a sequential postemergence treatment following a preemergence grass herbicide such as **Outlook**™, **Prowl**®, **Guardman Max**™, **Dual II Magnum**®, **Harness**®, or **Surpass**™. **IMPACT** may also be used in sequential programs with registered burn-down herbicides.

**IMPACT** may be applied in sequence with products containing isoxaflutole (e.g. **Balance**® Pro, **Epic**®, **Radius**™) if the isoxaflutole rate used is less than or equal to 0.0625 pounds active ingredient per acre (equal to 2 fluid ounces per acre of **Balance** Pro).

**IMPACT** may be applied in sequence with products containing mesotrione (e.g. **Callisto**®, **Lumax**®, **Camix**®, **Lexar**®) where corn will be planted the following growing season.

Pests listed in this label	
Common Name	Scientific Name
Amaranth, Palmer	<i>Amaranthus palmeri</i>
Amaranth, Powell	<i>Amaranthus powellii</i>
Barnyardgrass	<i>Echinochloa crus-galli</i>
Burcucumber	<i>Sicyos angulatus</i>
Carpetweed	<i>Mollugo verticillata</i>
Chickweed, Common	<i>Stellaria media</i>
Crabgrass, Large	<i>Digitaria sanguinalis</i>
Crabgrass, Smooth	<i>Digitaria ischaemum</i>
Cocklebur, Common	<i>Xanthium strumarium</i>
Cupgrass, Woolly	<i>Eriochloa villosa</i>
Dandelion	<i>Taraxacum officinale</i>
Foxtail, Giant	<i>Setaria faberi</i>
Foxtail, Green	<i>Setaria viridis</i>
Foxtail, Yellow	<i>Setaria glauca</i>
Galinsoga, Hairy	<i>Galinsoga ciliata</i>
Goosegrass	<i>Eleusine indica</i>
Horseweed (Marestail)	<i>Conyza canadensis</i>
Jimsonweed	<i>Datura stramonium</i>
Johnsongrass	<i>Sorghum halepense</i>
Kochia	<i>Kochia scoparia</i>
Lambsquarters, Common	<i>Chenopodium album</i>
Mallow, Common	<i>Malva neglecta</i>
Mallow, Venice	<i>Hibiscus trionum</i>
Millet, Wild-Proso	<i>Panicum miliaceum</i>
Morningglory	<i>Ipomoea sp.</i>
Mustard	<i>Brassica sp.</i>
Nightshade, Black	<i>Solanum nigrum</i>
Nightshade, Eastern Black	<i>Solanum ptycanthum</i>
Nightshade, Hairy	<i>Solanum sarrachoides</i>
Panicum, Fall	<i>Panicum dichotomiflorum</i>
Pigweed, Prostrate	<i>Amaranthus blitoides</i>
Pigweed, Redroot	<i>Amaranthus retroflexus</i>
Pigweed, Smooth	<i>Amaranthus hybridus</i>
Pigweed, Tumble	<i>Amaranthus album</i>
Prickly Lettuce	<i>Lactuca serriola</i>
Ragweed, Common	<i>Ambrosia artemisiifolia</i>
Ragweed, Giant	<i>Ambrosia trifida</i>
Shattercane	<i>Sorghum bicolor</i>
Sida, Prickly	<i>Sida spinosa</i>
Signalgrass, Broadleaf	<i>Brachiaria platyphylla</i>
Smartweed, Pennsylvania	<i>Polygonum pennsylvanicum</i>
Smartweed, Ladysthumb	<i>Polygonum persicaria</i>
Sunflower, Volunteer	<i>Helianthus sp.</i>
Sunflower, Wild (common)	<i>Helianthus annuus</i>
Thistle, Canada	<i>Cirsium arvense</i>
Thistle, Russian	<i>Salsola iberica</i>
Velvetleaf	<i>Abutilon theophrasti</i>
Waterhemp, Common	<i>Amaranthus rudis</i>
Waterhemp, Tall	<i>Amaranthus tuberculatus</i>

### LIMITED WARRANTY AND DISCLAIMER

The manufacturer warrants (a) that this product conforms to the chemical description on the label; (b) that this product is reasonably fit for the purposes set forth in the Directions for Use, subject to the inherent risks referred to herein, when it is used in accordance with such directions; and (c) that the directions, warnings, and other statements on this label are based upon responsible experts' evaluations of reasonable tests of effectiveness, of toxicity to laboratory animals and to plants and residues on food crops, and upon reports of field experience. Tests have not been made on all varieties of food crops and plants, or in all states or under all conditions.

**THERE ARE NO EXPRESS WARRANTIES OTHER THAN THOSE SET FORTH HEREIN. THE MANUFACTURER NEITHER MAKES NOR INTENDS, NOR DOES IT AUTHORIZE ANY AGENT OR REPRESENTATIVE, TO MAKE ANY OTHER WARRANTIES, EXPRESS OR IMPLIED, AND IT EXPRESSLY EXCLUDES AND DISCLAIMS ALL IMPLIED WARRANTIES OF MERCHANTABILITY OF FITNESS FOR A PARTICULAR PURPOSE, OR ANY WARRANTY OF QUALITY OR PERFORMANCE. THIS WARRANTY DOES NOT EXTEND TO, AND THE BUYER SHALL BE SOLELY RESPONSIBLE FOR, ANY AND ALL LOSS OR DAMAGE WHICH RESULTS FROM THE USE OF THIS PRODUCT IN ANY MANNER WHICH IS INCONSISTENT WITH THE LABEL DIRECTIONS, WARNINGS OR CAUTIONS.**

**BUYER'S EXCLUSIVE REMEDY AND MANUFACTURER'S OR SELLER'S EXCLUSIVE LIABILITY FOR ANY AND ALL CLAIMS, LOSSES, DAMAGES, OR INJURIES RESULTING FROM THE USE OR HANDLING OF THIS PRODUCT, WHETHER OR NOT BASED IN CONTRACT, NEGLIGENCE, STRICT LIABILITY IN TORT OR OTHERWISE, SHALL BE LIMITED, AT THE MANUFACTURER'S OPTION, TO REPLACEMENT OF, OR THE REPAYMENT OF THE PURCHASE PRICE FOR, THE QUANTITY OF PRODUCT WITH RESPECT TO WHICH DAMAGES ARE CLAIMED. TO THE FULLEST EXTENT PERMITTED BY LAW MANUFACTURER OR SELLER SHALL NOT BE LIABLE FOR SPECIAL, INDIRECT OR CONSEQUENTIAL DAMAGES RESULTING FROM THE USE OR HANDLING OF THIS PRODUCT.**

AMVAC offers this product, and Buyer accepts it, subject to the foregoing Limited Warranty which may be varied only by agreement in writing signed by an authorized representative of AMVAC.

IMPACT is a registered trademark of Amvac Chemical Corporation.

Clarity, Distinct, Lightning, Marksman, Prowl and Status are registered trademarks and G-Max Lite, Guardsman Max, and Outlook are trademarks of BASF Corporation.

Basagran, Clearfield and Laddok are trademarks of BASF AG.

Accent and Steadfast are registered trademarks and Cinch and Stout are trademarks of E.I. duPont de Nemours and Company.

Balance, Buctril, Epic, Liberty, LibertyLink, and Option are registered trademarks and Radius is a trademark of Bayer CropScience.

Callisto, Camix, Dual II Magnum, Exceed, Bicep II Magnum, Bicep Lite II Magnum, Lexar, Lumax, Spirit, Touchdown and Tough are registered trademarks of a Syngenta Group Company.

Degree, Degree Xtra, Harness, Harness Xtra, Permit, Roundup, Roundup Ready are registered trademarks of Monsanto Technology LLC.

Hornet, Keystone, Python, Stinger, Surpass, and TopNotch are trademarks of Dow AgroSciences LLC.

Aim is a registered trademark of FMC Corporation.

Amvac Chemical Corporation  
4100 E. Washington Boulevard  
Los Angeles, CA 90023